

Subject: Treatment Guidelines – BLS  
**Airway Adjunct Procedure**

Associated Policies:

---

- I. Indication
  - A. Decreased level of consciousness with absent or suppressed gag reflex.
- II. Contraindication
  - A. Absolute:
    - 1. Oral Airway:
      - a. Presence of a gag reflex.
    - 2. Nasal Airway:
      - a. Any head or facial trauma.
      - b. Child less than one (1) year of age.
- III. Procedure
  - A. Assemble equipment near patient:
    - 1. OPA/NPA.
    - 2. Oxygen.
    - 3. Bag valve mask.
    - 4. Suction.
  - B. Determine proper size:
    - 1. OPA:
      - a. Measure from the ear lobe to the corner of the mouth, or from the angle of the jaw to midline of the lips.
      - b. Do not use an undersized OPA, as there is risk of aspirating the airway.
    - 2. NPA:
      - a. Measure from the tip of the nostril to the tragus (cartilaginous bump) of the ear.
  - C. Insertion:
    - 1. OPA:
      - a. Gently insert the OPA upside down into the mouth, and advance the airway until resistance is met at the pharynx. When resistance is met, rotate the OPA 180° (right side up) while inserting remainder of airway or insert directly using tongue depressor to compress tongue.
    - 2. NPA:
      - a. Lubricate NPA with water soluble lubricant, and push up the tip of the nose to widen the nostrils. Insert into the larger or right nostril with the beveled side towards the septum. Angle the tip of the NPA at 45° above the base. Gently guide the NPA through the nostril while bringing it


Subject: Treatment Guidelines – BLS  
**Airway Adjunct Procedure**

---

to a 90° angle to the patient. Aim the tip of the NPA at C-1 and insert the remainder of the airway while rotating the airway back and forth between the fingertips. Do not attempt to force an NPA past resistance. Remove it and try the other nostril.

IV. Special Instructions

- A. Do not use an undersized OPA.
- B. Stimulation of an eyelash reflex is indicative of a gag reflex.
- C. Improper insertion of OPA's may push the tongue into the posterior pharynx and create an airway obstruction.
- D. Vigorous insertion of NPA's may cause severe nosebleeds.
- E. Never insert NPA's in each nostril.

Approved: 

Approved as to Form: 

Rev: 7/14